

The Interseasonal Resurgence of Respiratory Syncytial Virus in Australian Children Following the Reduction of Coronavirus Disease 2019–Related Public Health Measures

TO THE EDITOR—Yeoh et al reported the dramatic impact of public health measures introduced during the coronavirus disease 2019 (COVID-19) pandemic on influenza and respiratory syncytial virus (RSV) detections in Western Australian (WA) children [1]. Here, we present data from ongoing local prospective surveillance. Following the end of winter, there has been a persistent absence of severe acute respiratory syndrome coronavirus 2 community transmission and no increase in influenza detections. Limited

physical distancing measures have remained in place, with largely no restrictions on gathering sizes and no mandate for wearing masks [2]. Schools have remained open. Strict quarantine for overseas arrivals has been maintained, with a persistent marked decrease in visitor numbers compared with previous numbers [3]. Border restrictions for travelers from other states within Australia have been reduced as of 14 November 2020, with quarantine not required for travelers from states with no community severe acute respiratory syndrome coronavirus 2 transmission [4].

Similar to the previous report laboratory data prospectively collected as part of routine regional public health surveillance were collated per week from January 2012 to 13 December 2020. Cases were defined as detections of RSV

by validated nucleic acid amplification test or antigen detection kits in children (<16 years of age) in the metropolitan area. Laboratory results were provided by PathWest Laboratory Medicine, the only public pathology provider to the state. Samples were drawn from all public hospitals and emergency departments. Average epidemic curves for the period 2012 to 2019 were calculated using a World Health Organization–described method [5]. Median age was compared using 2-sample Wilcoxon test.

As demonstrated in Figure 1, RSV activity increased from late September, in the setting of relaxed physical distancing recommendations, ahead of the opening of interstate borders. Case numbers have further increased, exceeding the median seasonal peak from 2012 to 2019. This has been observed without any




Figure 1. Respiratory syncytial virus detection in children from metropolitan Western Australian up to week 50 of 2020, in the context of COVID-19 restrictions, compared to average epidemic curve (2012–2019). Borders closed – week 12 – international borders closed to all nonresidents; 14-day quarantine required for all arrivals. Period 1 (weeks 14–17): statewide stay-at-home restrictions in place with school holidays extended; borders closed to inter-state travelers (week 14). Period 2 (weeks 18–26): restrictions sequentially lifted, allowing gatherings of 10 (week 18), 20 (week 21), and 100 (week 24) people; schools reopened with increased cleaning and some physical distancing measures. Period 3 (weeks 27–47): majority of local restrictions removed, except the limitation of major sport and entertainment venues to 50% capacity; schools returned to all normal activities. Period 4 (weeks 47–50) relaxation of state border restrictions. Average epidemic curve: 2012–2019. Abbreviation: EYTD, end of year to date.

significant change in testing practices. The median patient age this year was 18.4 months, significantly higher than the upper range between 2012 and 2019 (7.3–12.5 months) ($P < .001$).

These data demonstrate the fragility of RSV control and the critical impact of physical distancing and respiratory hygiene practices. The rise in numbers and change in median age suggest that the expanded cohort of RSV-naïve patients, including an increased number of older children coupled with waning population immunity [6], may have contributed to this marked resurgence. Notably, RSV activity was first observed in mid-late August in regional centers before emergence in the metropolitan area, with transmission potentially facilitated by increased travel within the state [7]. Conversely, the initial rise in RSV cases in WA preceded the opening of interstate borders, suggesting this pathway was not the primary mechanism.

Our findings raise concerns for RSV control in the Northern Hemisphere, where a shortened season was experienced last winter [8]. The eventual reduction of COVID-19-related public health measures may herald a significant rise in RSV [9]. Depending on the timing, the accompanying morbidity and mortality, especially in older adults [6], may overburden already strained healthcare systems.

To delineate the relative contribution of an increasingly susceptible population and the reduction in border restrictions, further surveillance, transmission studies, and assessment of relatedness of RSV strains is progressing.

Notes

Financial support. This was an investigator-initiated project supported by a clinician-scientist

partnership grant from the Wesfarmers Centre for Vaccines and Infectious Diseases (2020).

D. K. Y. is supported by an Australian Government Research Training Program Postgraduate Scholarship. A. O. M. is supported by an NHMRC Postgraduate Scholarship (1191465) and an Australian Government Research Training Program Fees Offset. A. C. M. is supported by a Raine Clinician Research Fellowship. C. C. B. is supported by an NHMRC Career Development Fellowship (GNT1111596). H. C. M. is supported by a Telethon Kids Institute Emerging Research Leader Fellowship.

Potential conflicts of interest. A. O. M. reports grants from Australian Government (Research Training Program Fees Offset) and grants from NHMRC (Postgraduate Scholarship 1191465), outside the submitted work. A. C. M. reports grants from Raine Medical Research Foundation (Raine Clinician Research Fellowship), outside the submitted work. C. C. B. reports grants from NHMRC (Career Development Fellowship (GNT1111596)), outside the submitted work. D. K. Y. reports grants from Australian Government (Research Training Program Postgraduate Scholarship), outside the submitted work. H. C. M. reports from Telethon Kids Institute (Emerging Research Leader Fellowship) and grants (clinician-scientist partnership grant) and nonfinancial support from Wesfarmers Centre for Vaccines and Infectious Diseases, during the conduct of the study. All other authors: No reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

David A. Foley,^{1,*} Daniel K. Yeoh,^{1,2,*}
 Cara A. Minney-Smith,³ Andrew C. Martin,⁴
 Ariel O. Mace,^{4,5,6} Chisha T. Sikazwe,^{3,7}
 Huong Le,⁵ Avram Levy,^{3,7} Hannah C. Moore,⁸ and
 Christopher C. Blyth^{1,3,6,8}

¹Infectious Diseases Department, Perth Children's Hospital, Perth, Australia, ²Sir Peter MacCallum Department of Oncology, University of Melbourne, Parkville, Victoria, ³Department of Microbiology, PathWest Laboratory Medicine WA, Perth, Australia, ⁴Department of General Paediatrics, Perth Children's Hospital, Perth, Australia, ⁵Department of General Paediatrics, Fiona Stanley Hospital, Perth, Australia, ⁶Wesfarmers Centre for Vaccines and Infectious Diseases, Telethon Kids Institute, University of Western Australia, Perth, Australia, ⁷Faculty of Health and Medical Sciences, University of Western Australia, Perth, Australia, and ⁸School of Medicine, University of Western Australia, Perth, Australia

References

- Yeoh DK, Foley DA, Minney-Smith CA, et al. The impact of COVID-19 public health measures on detections of influenza and respiratory syncytial virus in children during the 2020 Australian winter. *Clin Infect Dis* 2020.
- COVID-19 coronavirus: WA roadmap [press release]. Western Australia Government Website. 2020. Available at: <https://www.wa.gov.au/organisation/department-of-the-premier-and-cabinet/covid-19-coronavirus-wa-roadmap>. Accessed 15 December 2020.
- Australian Bureau of Statistics. Overseas arrivals and departures Australia 2020. Available at: <https://www.abs.gov.au/statistics/industry/tourism-and-transport/overseas-arrivals-and-departures-australia/latest-release>. Accessed 6 December 2020.
- Department of the Premier and Cabinet. Covid 19 Coronavirus: what you can and can't do 2020. Available at: <https://www.wa.gov.au/organisation/department-of-the-premier-and-cabinet/covid-19-coronavirus-what-you-can-and-cant-do>. Accessed 6 December 2020.
- World Health Organization. Global epidemiological surveillance standards for influenza. 2013. Available at: https://www.who.int/influenza/resources/documents/WHO_Epidemiological_Influenza_Surveillance_Standards_2014.pdf?ua=1. Accessed 1 October 2020.
- Lambert L, Sagfors AM, Openshaw PJ, Culley FJ. Immunity to RSV in early-life. *Front Immunol* 2014; 5:466.
- Apple. Mobility trend reports 2020. Available at: <https://covid19.apple.com/mobility>. Accessed 8 December 2020.
- Kuitunen I, Artama M, Mäkelä L, Backman K, Heiskanen-Kosma T, Renko M. Effect of social distancing due to the COVID-19 pandemic on the incidence of viral respiratory tract infections in children in finland during early 2020. *Pediatr Infect Dis J* 2020; 39:e423–7.
- Baker RE, Park SW, Yang W, Vecchi GA, Metcalf CJE, Grenfell BT. The impact of COVID-19 nonpharmaceutical interventions on the future dynamics of endemic infections. *Proc Natl Acad Sci U S A* 2020; 117:30547–53.

*D.A.F. and D.K.Y. contributed equally to this manuscript. Correspondence: D. A. Foley, Department of Infectious Diseases, Perth Children's Hospital, 15 Hospital Ave, Perth, Western Australia 6009, Australia (drdavidanthonyfoley@gmail.com).

Clinical Infectious Diseases® 2021

© The Author(s) 2021. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com. DOI: 10.1093/cid/ciaa1906